

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 February 2001 (01.02.2001)

PCT

(10) International Publication Number
WO 01/07625 A2

(51) International Patent Classification⁷: C12N 15/31, C07K 14/29, C12N 9/52, 9/02, A61K 39/00

(21) International Application Number: PCT/US00/19763

(22) International Filing Date: 20 July 2000 (20.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/358,322 21 July 1999 (21.07.1999) US

(71) Applicant: CORNELL RESEARCH FOUNDATION, INC. [US/US]; 20 Thornwood Drive, Suite 105, Ithaca, NY 14850 (US).

(72) Inventor: CHANG, Yung-fu; 204 Christopher Lane, Ithaca, NY 14850 (US).

(74) Agents: BROWN, Michael, F. et al.; Brown, Pinnisi & Michaels, P.C., Suite 400, 118 North Tioga Street, Ithaca, NY 14850 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: EHRLICHIA CANIS GENES AND VACCINES



(57) Abstract: This invention provides the sequence of 5,300 nucleotides from the E. canis genome. There are four proteins, ProA, ProB, ORF, and a cytochrome oxidase homolog, as well as a partial lipoprotein signal peptidase homolog at the carboxy terminus, coded for in this cloned fragment. The antigenic properties of these proteins allow them to be used to create a vaccine. An embodiment of this invention includes the creation of a DNA vaccine, a recombinant vaccine, and a T cell epitope vaccine.

***Ehrlichia canis* Genes and Vaccines**

FIELD OF THE INVENTION

The invention pertains to the field of veterinary pathogens. More particularly, the present invention pertains to the sequence of specific genes of the bacterial canine pathogen *Ehrlichia canis* and the application of this technology to the development of a vaccine.

BACKGROUND OF THE INVENTION

The present invention relates to the sequence of genes from the *E. canis* bacterium, and the development of a vaccine against this organism.

Ehrlichia canis (*E. canis*) is a small gram-negative, obligately intracellular bacterium. This bacteria is the agent which causes canine monocytic ehrlichiosis (CME), a tick-borne disease which predominantly affects dogs. The most common carrier of *E. canis* is the brown dog tick *Rhipicephalus sanguineus*. The disease was described originally in Algeria in 1935. It was subsequently recognized in the United States in 1962, but is now known throughout much of the world. Canine monocytic ehrlichiosis caused much concern during the Vietnam War, when 160 military dogs died from the *E. canis* infection. There is no vaccination currently available against *E. canis*. It is a life threatening disease that continues to be an important health concern for veterinarians and pet owners alike.

Canine monocytic ehrlichiosis is an infectious blood disease. A reduction in cellular blood elements is the primary characteristic of the disease. *E. canis* lives and reproduces in the white blood cells (leukocytes). It eventually affects the entire lymphatic system, and devastates multiple organs. By targeting the white blood cells, these cells die

off rapidly. These dead blood cells migrate primarily to the spleen, which enlarges as a result. The bone marrow recognizes the loss of the white blood cells and works to form new, healthy cells. It sends out the cells prematurely, and these immature cells do not work properly. Often, these immature cells mimic those in leukemic patients, so the disease is misdiagnosed as leukemia. Canine monocytic ehrlichiosis may predispose dogs to various cancers.

There are three stages of canine monocytic ehrlichiosis. The first, acute stage mimics a mild viral infection. During the acute stage, most, if not all, of the damage is reversible and the animal is likely to recover. This is the stage where treatment is the most effective, stressing the need for early detection. Without treatment, however, the animal will progress into a subclinical (second) stage and/or to the chronic (final) stage. When the animal has reached the chronic stage, the bacterial organism has settled within the bone marrow. Many dogs in this stage suffer massive internal hemorrhage, or develop lethal complications such as sudden stroke, heart attack, renal failure, splenic rupture or liver failure.

E. canis can be cultured *in vitro* in a mammalian-derived cell line (DH82). Continued maintenance of these cells is difficult because the cell culture must be supplemented with primary monocytes (white blood cells found in bone marrow) every two weeks. The cultures are very slow growing, and the culture media is expensive.

Data concerning the genes in the *E. canis* genome has concentrated primarily on the 16S rRNA gene. Previous work has sequenced this gene, which is a ubiquitous component of the members of the ehrlichia family, as well as the majority of organisms worldwide. The high sequence homology between this gene throughout the living world makes it a poor candidate for vaccine development. It is necessary to find other genes within this genome if hope for a vaccine against this deadly disease can ever be realized.

Sequencing of the 16S rRNA gene indicates that *E. canis* is closely related (98.2% homology) to *E. chaffeensis*, the novel etiologic agent of human ehrlichiosis. Western blots of *E. canis* are similar when probed with antisera to *E. canis*, *E. chaffeensis* and *E.*

ewingi (another cause of human ehrlichiosis) indicating a close antigenic relationship between these three species (Chen *et al.*, 1994).

The indirect fluorescent antibody test (IFA) has been developed for detecting canine monocytic ehrlichiosis. IFA detects the presence of antibodies against the invading organism in a dog's blood. Unfortunately, this test is not always accurate. Sometimes, dogs will test negative in the acute phase because their immune system is delayed in forming antibodies. Another false negative may occur if there is a low titer in the chronic stage. An additional drawback of this test is the cross-reactivity found. The anti *E. canis* polyclonal antibody positively reacts with *E. chaffeensis*, undermining the specificity of the test. An alternative test, the Giesma smear, has been used to locate the actual organism in a dog's blood. Unfortunately, despite appropriate staining techniques and intensive film examination, the organisms frequently can not be located. The fallibility of these tests makes it essential to provide better diagnostic tools for this disease.

Due to difficulties in the detection of a tick bite, early diagnosis of infection, the suppression of host defenses and the nature of persistent infection of the disease, an effective vaccine against *E. canis* is urgently needed for dogs.

SUMMARY OF THE INVENTION

This invention discloses novel sequence data for *E. canis* genes. Specifically, a clone has been identified and sequenced. Four proteins termed ProA, ProB, ORF (an open reading frame with unknown function) and a cytochrome oxidase homolog, have been identified within this clone. In addition, a partial gene encoding a lipoprotein signal peptidase homolog has been discovered.

An embodiment of this invention includes the creation of a vaccine with this sequence and protein information. The proteins disclosed in this invention are extremely antigenic. Therefore, they have the potential to be extremely useful as a vaccine. The

types of vaccine made available by this novel technology include a DNA vaccine, a recombinant vaccine, and a T cell epitope vaccine.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the three clones identified in the library screen.

DESCRIPTION OF THE PREFERRED EMBODIMENT

E. canis causes a devastating canine disease. Currently, there is no vaccine available to prevent this disease. This invention provides the tools necessary to develop such a vaccine. More specifically, four genes have been identified from a genomic fragment of *E. canis*, named ProA, ProB, ORF and a cytochrome oxidase homolog. In addition, a partial gene coding for a lipoprotein signal peptidase homolog has been found. Any of these proteins can be utilized in an embodiment of this invention to develop a vaccine.

Screening an *E. canis* library

To identify genes in the *E. canis* genome, a genomic DNA expression library was constructed. An *E. canis* strain isolated from dogs with canine ehrlichiosis was grown in the dog cell line DH82 by a technique being known in the art, and incorporated by reference (Dawson *et al.*, 1991; Rikihisa, 1992). The cells were harvested and the chromosomal DNA extracted as described by a technique known in the art (Chang *et al.*, 1987; Chang *et al.*, 1989a; Chang *et al.*, 1989b; Chang *et al.*, 1993a; Chang *et al.*, 1993b). To construct the library, 200 µg of DNA was partially digested with *Sau*3A. DNA fragments from 3 to 8 kb were isolated and ligated to a plasmid, pHG165 (Stewart *et al.*, 1986). The plasmids were transformed into *E. coli* TB1 (Chang *et al.*, 1987).

The library was screened with polyclonal antibodies against *E. canis*. Polyclonal antibodies were generated from dogs that had been bitten by a tick harboring *E. canis*.

The polyclonal antibodies were preabsorbed with the lysate of an *E. coli* host strain. The library was plated on petri plates at a density of 1,000 colony forming units. Colonies were transferred to nitrocellulose and each filter was probed with 1 ml of the preabsorbed polyclonal antibodies. Positive colonies were identified with a second antibody consisting of an alkaline phosphatase-conjugated goat anti-rabbit IgG (Kirkegaard and Perry Laboratories, Gaithersburg, MD), followed by color development with a substrate solution containing nitroblue tetrazolium (NBT) and 5-bromo-4-chloro-3-indolyl phosphate (BCIP). Positive clones were rescreened three times.

Three clones were isolated from this screening procedure (Figure 1). The longest genomic fragment (pCH4) encodes four complete genes and one partial gene. It completely encodes the proteins ProA, ProB, ORF and a cytochrome oxidase homolog, as well as containing the partial sequence of a lipoprotein signal peptidase homolog. ProA and ProB are located on a single operon. Restriction endonuclease digestion mapping and DNA sequencing were done by techniques known in the art, and incorporated by reference (Chang *et al.*, 1987; Chang *et al.*, 1989a; Chang *et al.*, 1989b; Chang *et al.*, 1993a; Chang *et al.*, 1993b). Briefly, the DNA sequence was determined by automated DNA sequencing on the ABI PRISM Model 377 DNA system. The complete nucleotide sequences were determined on both strands by primer walking. The thermal cycling of the sequencing reactions utilized the Taq DyeDeoxyTM Terminator Cycle sequencing kit. Databases were searched for homologous proteins through the use of the BLAST network service of the National Center for Biotechnology Information (NCBI) (Althchul *et al.*, 1990; Gish *et al.*, 1993).

Sequence Information

The *E. canis* genes were sequenced. The cloned fragment contains 5,300 nucleotides, and codes for four proteins. There is also one partial gene at the carboxy terminus. SEQ. ID. NO. 1 is the entire nucleotide sequence. SEQ. ID. NO. 2 and 3 are the translation of nucleotides 12 through 533 from SEQ. ID. NO. 1 and code for a cytochrome oxidase homolog. Cytochrome oxidase is important in virulence, and therefore is a strong candidate for use in a vaccine. SEQ. ID. NO. 4 and 5 are the translation of nucleotides 939 through 2,252 from SEQ. ID. NO. 1 and code for ProA. SEQ. ID. NO. 6 and 7 are the

translation of nucleotides 2,258 through 3,664 from SEQ. ID. NO. 1 and code for ProB. Preliminary evidence indicates that ProA and ProB are proteases. SEQ. ID. NO. 8 and 9 are the translation of nucleotides 4,121 through 4,795 from SEQ. ID. NO. 1 and code for ORF, a protein with unknown function. SEQ. ID. NO. 10 and 11 are the translation of the complementary sequence of nucleotides 4,884 through 5,300 from SEQ. ID. NO. 1 and code for the partial sequence of a lipoprotein signal peptidase homolog. Lipoprotein signal peptidases are membrane proteins, and by nature may be less desirable for vaccine development. However, this protein is still worth pursuing in the creation of a vaccine.

Overexpression of ProA, ProB, ORF, cytochrome oxidase and the lipoprotein signal peptidase homolog

The *E. canis* antigens are overexpressed in a T7 promoter plasmid. The pRSET vector allows high level expression in *E. coli* in the presence of T7 RNA polymerase, which has a strong affinity for the T7 promoter. After subcloning the antigen genes into the pRSET vector, the subclones are transformed into an F' *E. coli* JM109 strain. For maximum protein expression, the transformants are cultured to O.D. 600=0.3, exposed to IPTG (1 mM) for one hour and then transfected with M13/T7 bacteriophages at a multiplicity of infection (MOI) of 5-10 plaque forming units (pfu) per cell. Time course studies indicate that maximum induction is reached two hours after induction.

The pellet is harvested by centrifugation and the cells are resuspended in 6M Guanidinium (pH 7.8). Cells are ruptured by French press and the total lysate is spun at 6000 rpm to separate cell debris by a technique known in the art, and hereby incorporated by reference (Chang *et al.*, 1993c). Immobilized metal ion affinity chromatography (IMIAc) is used to purify each of the proteins under denaturing conditions as described by the manufacturer (Invitrogen, San Diego, CA). The protein samples are separated by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and visualized after staining with coomassie blue.

Vaccine Development

Prior to the present invention, no vaccine against *E. canis* had been developed. *E. canis* is endemic in dogs and closely related canidae in many parts of the world. Dogs in

North America are also increasingly at risk and the application of the present invention can potentially save the lives of thousands of dogs each year. An *E. canis* vaccine that can elicit cell-mediated immunity against this tick-borne disease of dogs is desperately needed.

DNA Vaccine

A DNA vaccine is constructed by subcloning the gene of interest into a eukaryotic plasmid vector. Candidate vectors include, but are not limited to, pcDNA3, pCI, VR1012, and VR1020. This construct is used as a vaccine.

Each of the newly identified genes, ProA, ProB, ORF, the cytochrome oxidase homolog, or the partial lipoprotein signal peptidase homolog can be used to create a DNA vaccine (reviewed in Robinson, 1997). In addition, any immunologically active portion of these proteins is a potential candidate for the vaccine. A plasmid containing one of these genes in an expression vector is constructed. The gene must be inserted in the correct orientation in order for the genes to be expressed under the control of eukaryotic promoters. Possible promoters include, but are not limited to, the cytomegalovirus (CMV) immediate early promoter, the human tissue plasminogen activator (t-PA) gene (characterized in Degen *et al.*, 1986), and the promoter/enhancer region of the human elongation factor alpha (EF-1 α) (characterized in Uetsuki *et al.*, 1989). Orientation is identified by restriction endonuclease digestion and DNA sequencing.

Expression of these gene products is confirmed by indirect immunofluorescent staining of transiently transfected COS cells. The same plasmid without these genes is used as a control. Plasmid DNA is transformed into *Escherichia coli* DH5 α . DNA is purified by cesium chloride gradients and the concentration is determined by a standard protocol being known in the art, and incorporated by reference (Nyika *et al.*, 1998).

Once the vector is purified, the vector containing the DNA can be suspended in phosphate buffer saline solution and directly injected into dogs. Inoculation can be done via the muscle with a needle or intravenously. Alternatively, a gene gun can be used to transport DNA-coated gold beads into cells by a technique known in the art, and hereby incorporated by reference (Fynan *et al.*, 1993). The rationale behind this type of vaccine

is that the inoculated host expresses the plasmid DNA in its cells, and produces a protein that raises an immune response. Each of the newly identified genes can be used to create a vaccine by this technique.

CpG molecules can be used as an adjuvant in the vaccine. This technique is known in the art, and is hereby incorporated by reference (Klinman *et al.*, 1997). Adjuvants are materials that help antigens or increase the immune response to an antigen. The motifs consist of an unmethylated CpG dinucleotide flanked by two 5' purines and two 3' pyrimidines. Oligonucleotides containing CpG motifs have been shown to activate the immune system, thereby boosting an antigen-specific immune response. This effect can be utilized in this invention by mixing the CpG oligonucleotides with the DNA vaccine, or physically linking the CpG motifs to the plasmid DNA.

Recombinant Vaccine

In order to develop a recombinant vaccine, each of the genes is individually subcloned into overexpression vectors, and then purified for vaccine development. ProA, ProB, ORF, the cytochrome oxidase homolog or the partial lipoprotein signal peptidase homolog is expressed in a plasmid with a strong promoter such as the tac, T5, or T7 promoter. Alternatively, immunologically active fragments of these proteins are used in the development of a vaccine. Each of these genes is subcloned into a plasmid and transformed into an *E. coli* strain as described above.

The recombinant protein is overexpressed using a vector with a strong promoter. Vectors for use in this technique include pREST (Invitrogen Inc., CA), pKK233-3 (Pharmacia, CA), and the pET system (Promega, WI), although any vector with a strong promoter can be used. After overexpression, the proteins are purified and mixed with adjuvant. Potential adjuvants include, but are not limited to, aluminum hydroxide, QuilA, or Montamide. The purified protein is used as immunogen to vaccinate dogs by a technique being known in the art, and incorporated by reference (Chang *et al.*, 1993c; Chang *et al.*, 1995). Briefly, the individual protein is expressed and purified from *E. coli*. Then, the dogs are injected intramuscularly or subcutaneously with the purified recombinant vaccine and adjuvant. This injection elicits an immune response.

T Cell Epitope Vaccine

Direct cell cytotoxicity mediated by CD8⁺ T lymphocytes (CTL) is the major mechanism of defense against intracellular pathogens. These effector lymphocytes eliminate infected cells by recognizing short peptides associated with MHC class I molecules on the cell surface. Exogenous antigens enter the endosomal pathway and are presented to CD4⁺ T cells in association with class II molecules whereas endogenously synthesized antigens are presented to CD8⁺ T cells in association with MHC class I molecules. *E. canis* is an intracellular pathogen that resides in monocytes and macrophages. The present invention develops novel ways of generating an *E. canis*-specific CTL response that would eliminate the organism from monocytes or macrophages of infected animals.

A strategy for increasing the protective response of a protein vaccine is to immunize with selective epitopes of the protein. The rationale behind this is that an epitope vaccine contains the most relevant immunogenic peptide components without the irrelevant portions. Therefore, a search is performed for the most highly antigenic portions of the newly identified proteins.

To identify T-cell epitopes from the newly discovered proteins, an initial electronic search for homologous sequences to known T-cell epitopes is performed. In addition, extensive T-cell epitope mapping is carried out. Each of the proteins, ProA, ProB, ORF, the cytochrome oxidase homolog, and the partial lipoprotein signal peptidase homolog, is tested for immunogenic peptide fragments. Mapping of T cell epitopes by a technique known in the art is hereby incorporated by reference (Launois *et al.*, 1994; Lee and Horwitz, 1999). Briefly, short, overlapping peptide sequences (9-20 amino acids) are synthesized over the entire length of the protein in question. These short peptide fragments are tested using healthy dogs which have been immunized with the protein of interest. Peripheral blood mononuclear cells from the dogs are tested for T cell stimulatory and IFN- γ inducing properties. Those fragments which elicit the strongest response are the best candidates for a T-cell epitope vaccine.

Once fragments are identified which will make the best epitopes, a recombinant adenylate cyclase of *Bordetella bronchiseptica* is constructed carrying an *E. canis* CD8⁺ T cell epitope. The adenylate cyclase toxin (CyaA) of *Bordetella bronchiseptica* causes disease in dogs and elicits an immune response. In addition, CyaA is well suited for intracytoplasmic targeting. Its catalytic domain (AC), corresponding to the N-terminal 400 amino acid residues of the 1,706-residue-long protein, can be delivered to many eukaryotic cells, including cells of the immune system. Also, toxin internalization is independent of receptor-mediated endocytosis, suggesting that the catalytic domain can be delivered directly to the cytosol of target cells through the cytoplasmic membrane. The *Pseudomonas aeruginosa* exotoxin A (PE) is another toxin which could be used in this procedure to deliver peptides or proteins into cells, by a technique known in the art, and hereby incorporated by reference (Donnelly *et al.*, 1993).

Foreign peptides (16 residues) have been inserted into various sites of the AC domain of CyaA without altering its stability or catalytic and calmodulin-binding properties. Thus, protein engineering allows the design and delivery of antigens that specifically stimulate CTLs. The induction of specific CD8⁺ T cells can play an important role in canine ehrlichiosis control due to the intracellular persistence of *E. canis* in monocytes.

The adenylate cyclase (AC) toxin (*cya*) gene of *B. bronchiseptica* has been cloned. A synthetic double-stranded oligonucleotide encoding a 9 to 20 amino acid class I T cell epitope of either ProA, ProB, ORF, the cytochrome oxidase homolog, or the partial lipoprotein signal peptidase homolog, is designed according to *B. bronchiseptica* codon usage. The complementary oligonucleotides are inserted in the hypervariable region of the cloned AC-coding sequence of the *cya*. This technique is known in the art in other systems, and is incorporated by reference (Sebo *et al.*, 1995; Guernonprez *et al.*, 1999).

Recombinant plasmids carrying the chimeric *cya* gene are sequenced to determine the copy number and orientation of the inserted epitope. A plasmid with a complete copy of the insert that specifies the T-cell epitope (CD8⁺) in the correct orientation is chosen from the sequenced plasmids. The ability of the new chimeric protein to enter eukaryotic cells is necessary to ensure intracellular targeting of the epitopes (Fayolle *et al.*, 1996).

A vaccine can be created in one of two ways. Recombinant chimeric protein can be purified and used to inoculate dogs. Alternatively, an attenuated *B. bronchiseptica* strain that carries a T-cell epitope or *E. canis* gene by in-frame insertion into adenylate cyclase is created by allelic-exchange. Allelic-exchange is a technique known in the art, and is hereby incorporated by reference (Cotter and Miller, 1994).

Finally, protection against *E. canis* infection in dogs vaccinated with the adenylase cyclase- ProA, ProB, ORF, cytochrome oxidase homolog, or lipoprotein signal peptidase homolog chimeric protein is determined. Wild type and recombinant ACs and CyAs are diluted to working concentrations in PBS and the chimeric protein is injected into dogs either intramuscularly or subcutaneously. Alternatively, the T-cell epitope is inserted into the adenylate cyclase gene of an attenuated *B. bronchiseptica* strain in frame, and the dogs are given the live bacteria.

Recombinant antigens are promising candidates for human and animal vaccination against various pathogens. However, a serious drawback is the poor immunogenicity of recombinant antigens as compared to native antigens. A major challenge in the development of a new recombinant vaccine is, therefore, to have a new adjuvant system that increases the immunogenicity of antigens. Cytokines are powerful immunoregulatory molecules. Cytokines which could be used as adjuvants in this invention include, but are not limited to, IL-12 (interleukin-12), GM-CSF (granulocyte-macrophage colony stimulating factor), IL-1 β (interleukin-1 β) and γ -IFN (gamma interferon).

These cytokines can have negative side effects including pyrogenic and/or proinflammatory symptoms in the vaccinated host. Therefore, to avoid the side effects of a whole cytokine protein, an alternate approach is to use synthetic peptide fragments with the desired immunostimulatory properties. The nonapeptide sequence VQGEESNDK of IL-1 β protein is endowed with powerful immuno-enhancing properties, and is discussed here to illustrate the use of a cytokine to increase immunogenicity.

This nonapeptide is inserted into the ProA, ProB, ORF, the cytochrome oxidase homolog, or the partial lipoprotein signal peptidase homolog protein and its immunogenicity is compared to that of the native protein. Reportedly, the insertion of this

sequence into a poorly immunogenic recombinant antigen increases the chance of a strong protective immune response after vaccination. This peptide could enhance the *in vivo* immune response against both T-dependent and T-independent antigens. The canine IL-1 β sequence may mimic many immunomodulatory activities of the entire molecule of IL-1 β while apparently lacking many of its undesirable proinflammatory properties. This strategy is employed to increase the immunogenicity of ProA, ProB, ORF, cytochrome oxidase, the partial lipoprotein signal peptidase homolog and other *E. canis* antigens.

Plasmid pYFC199 is derived from a pBR322 plasmid by the insertion of a fragment that includes the ProA, ProB, ORF, the cytochrome oxidase homolog, or the partial lipoprotein signal peptidase protein from *E. canis*. This plasmid contains a unique *Hind*III site where in-frame insertions encoding exogenous sequences can be inserted. Two complementary oligonucleotides, AGGCTTGTTTCAGGGTGAAGAAGAATCCAACGACAAAAGCTT and AAGCTTTTGTCGTTGGATTCTTCACCCTGAAGTTGCCA, that encode the canine IL-1 β 163-171 peptide are annealed, cut with *Hind*III, and inserted into the pYFC199 *Hind*III site. The recombinant plasmid carrying the chimeric IL-1 β gene is sequenced to determine the orientation of the inserted epitope.

The efficacy of the recombinant proteins as vaccines is tested in dogs. The purified protein is injected intraperitoneally into dogs. Specific pathogen free (SPF) dogs are divided into five groups: one group is given recombinant adenylate cyclase of *Bordetella bronchiseptica* carrying *E. canis* CD8⁺ T cell epitopes derived from ProA, ProB, ORF, cytochrome oxidase homolog, or the partial lipoprotein signal peptidase homolog, one group is given recombinant adenylate cyclase of *Bordetella bronchiseptica* as a control, one group is given the ProA, ProB, ORF, cytochrome oxidase homolog, or the partial lipoprotein signal peptidase homolog protein plus a canine IL-1 β 163-171 insert, one group is given a T cell epitope derived from ProA, ProB, ORF, cytochrome oxidase homolog, or the partial lipoprotein signal peptidase homolog alone, and the last group is given PBS as a negative control.

All animals are vaccinated (30-40 μ g each) four times. The dogs are challenged ten days after the last vaccination with 10^7 *E. canis*. At day five postchallenge, approximately 1 ml blood from each dog is collected in an EDTA tube. Whether the vaccinated groups eliminate the organisms as compared to that of the control group is tested by culture and PCR. Two primers derived from the genes cloned can be used to amplify the gene product from the tissues or blood samples from these dogs. The internal primer can also be designed for use as an oligonucleotide probe to hybridize the PCR gene product.

This invention provides a badly needed vaccine against the *E. canis* bacterium. The vaccine can be used to protect dogs throughout the world from canine monocytic ehrlichiosis.

Accordingly, it is to be understood that the embodiments of the invention herein described are merely illustrative of the application of the principles of the invention. Reference herein to details of the illustrated embodiments are not intended to limit the scope of the claims, which themselves recite those features regarded as essential to the invention.

What is claimed is:

- 1 1. A recombinant DNA comprising said DNA selected from the group consisting of:
 - 2 a) a recombinant DNA that encodes a protein having an amino acid sequence as
3 shown in SEQ. ID. NO. 3;
 - 4 b) a recombinant DNA that encodes a protein having an amino acid sequence as
5 shown in SEQ. ID. NO. 5;
 - 6 c) a recombinant DNA that encodes a protein having an amino acid sequence as
7 shown in SEQ. ID. NO. 7;
 - 8 d) a recombinant DNA that encodes a protein having an amino acid sequence as
9 shown in SEQ. ID. NO. 9;
 - 10 e) a recombinant DNA that encodes a protein having an amino acid sequence as
11 shown in SEQ. ID. NO. 11; and
 - 12 f) any portion of said DNA above that encodes a protein that elicits an immune
13 response against *E. canis*.
- 1 2. The recombinant DNA of claim 1 wherein said DNA encodes at least one
2 immunogenic epitope.
- 1 3. A recombinant protein comprising said protein selected from the group consisting of:
 - 2 a) a protein having an amino acid sequence as shown in SEQ. ID. NO. 3;
 - 3 b) a protein having an amino acid sequence as shown in SEQ. ID. NO. 5;
 - 4 c) a protein having an amino acid sequence as shown in SEQ. ID. NO. 7;
 - 5 d) a protein having an amino acid sequence as shown in SEQ. ID. NO. 9;
 - 6 e) a protein having an amino acid sequence as shown in SEQ. ID. NO. 11; and

7 f) any portion of any of the above proteins that elicits an immune response
8 against *E. canis*.

1 4. The recombinant protein of claim 3 wherein said protein includes at least one
2 immunogenic epitope.

1 5. A vaccine wherein said vaccine protects dogs against *E. canis* infection.

1 6. The vaccine of claim 5 comprising:

2 a) a vector capable of expressing a recombinant DNA inserted into said vector
3 such that a recombinant protein is expressed when said vector is provided in an
4 appropriate host; and

5 b) the recombinant DNA inserted into said vector wherein said DNA is selected
6 from the group consisting of:

7 i. a recombinant DNA that encodes a protein having an amino acid
8 sequence as shown in SEQ. ID. NO. 3;

9 ii. a recombinant DNA that encodes a protein having an amino acid
10 sequence as shown in SEQ. ID. NO. 5;

11 iii. a recombinant DNA that encodes a protein having an amino acid
12 sequence as shown in SEQ. ID. NO. 7;

13 iv. a recombinant DNA that encodes a protein having an amino acid
14 sequence as shown in SEQ. ID. NO. 9;

15 v. a recombinant DNA that encodes a protein having an amino acid
16 sequence as shown in SEQ. ID. NO. 11; and

17 vi. any portion of said DNA above that encodes a protein that elicits an
18 immune response against *E. canis*.

1 7. The vaccine of claim 6, wherein said DNA further comprises DNA that encodes CpG
2 motifs.

1 8. The vaccine of claim 6 wherein said DNA further comprises a promoter selected from
2 the group consisting of:

3 a) a cytomegalovirus (CMV) immediate early promoter;

4 b) a human tissue plasminogen activator gene (t-PA); and

5 c) a promoter/enhancer region of a human elongation factor alpha (EF-1 α).

1 9. The vaccine of claim 6, wherein said vector is selected from the group consisting of:

2 a) pcDNA3;

3 b) pC1;

4 c) VR1012; and

5 d) VR1020.

1 10. The vaccine of claim 6 wherein said vaccine is administered into said host by a method
2 selected from the group consisting of:

3 a) intramuscular injection;

4 b) intravenous injection; and

5 c) gene gun injection.

1 11. The vaccine of claim 10, wherein said host is a dog.

1 12. The vaccine of claim 5 comprising:

2 a) a recombinant protein that is selected from the group consisting of:

3 i. a protein having an amino acid sequence as shown in SEQ. ID. NO. 3;

- 4 ii. a protein having an amino acid sequence as shown in SEQ. ID. NO. 5;
5 iii. a protein having an amino acid sequence as shown in SEQ. ID. NO. 7;
6 iv. a protein having an amino acid sequence as shown in SEQ. ID. NO. 9;
7 v. a protein having an amino acid sequence as shown in SEQ. ID. NO. 11;
8 and
9 vi. any portion of any of the above proteins that elicits an immune response
10 against *E. canis*.

1 13. The vaccine of claim 12, wherein said vaccine further comprises adjuvants selected
2 from the group consisting of:

- 3 a) aluminum hydroxide;
4 b) QuilA; and
5 c) Montamide.

1 14. The vaccine of claim 12 further comprising a cytokine operatively associated with said
2 recombinant protein.

1 15. The vaccine of claim 14 wherein said cytokine is selected from the group consisting
2 of:

- 3 a) interleukin-1 β (IL-1 β);
4 b) granulocyte-macrophage colony stimulating factor (GM-CSF);
5 c) gamma interferon (γ -IFN);
6 d) amino acids VQGEESENDK from the IL-1 β protein; and
7 e) any portion of any of the cytokines above that elicits an improved
8 immunogenic response against *E. canis*.

1 16. The vaccine of claim 12 wherein said vaccine is administered into a host by a method
2 selected from the group consisting of:

- 3 a) intramuscular injection; and
4 b) subcutaneous injection.

1 17. The vaccine of claim 16 wherein said host is a dog.

1 18. The vaccine of claim 5 comprising a recombinant protein that includes a T cell epitope
2 wherein said T cell epitope comprises an amino acid peptide fragment of a protein
3 selected from the group consisting of:

- 4 a) a protein having an amino acid sequence as shown in SEQ. ID. NO. 3;
5 b) a protein having an amino acid sequence as shown in SEQ. ID. NO. 5;
6 c) a protein having an amino acid sequence as shown in SEQ. ID. NO. 7;
7 d) a protein having an amino acid sequence as shown in SEQ. ID. NO. 9;
8 e) a protein having an amino acid sequence as shown in SEQ. ID. NO. 11; and
9 f) any portion of any of the above proteins that elicits an immune response
10 against *E. canis*.

1 19. The vaccine of claim 18 wherein said amino acid peptide fragment comprises nine to
2 twenty amino acids.

1 20. The vaccine of claim 18 further comprising a recombinant DNA encoding a protein
2 which is capable of being internalized into eukaryotic cells, including cells of the
3 immune system.

1 21. The vaccine of claim 20 wherein said protein capable of being internalized into
2 eukaryotic cells comprises a toxin selected from the group consisting of:

- 3 a) a recombinant adenylate cyclase of *Bordetella bronchiseptica*; and

4 b) a recombinant exotoxin A (PE) of *Pseudomonas aeruginosa*.

1 22. The vaccine of claim 18 wherein said vaccine is administered into a host by a method
2 selected from the group consisting of:

3 a) intramuscular injection; and

4 b) subcutaneous injection.

1 23. The vaccine of claim 22 wherein said host is a dog.

1 24. A method of identifying a T cell epitope against *E. canis* comprising:

2 a) synthesizing overlapping peptide fragments over an entire length of a protein
3 wherein said protein is selected from the group consisting of:

4 i. a protein having an amino acid sequence as shown in SEQ. ID. NO. 3;

5 ii. a protein having an amino acid sequence as shown in SEQ. ID. NO. 5;

6 iii. a protein having an amino acid sequence as shown in SEQ. ID. NO. 7;

7 iv. a protein having an amino acid sequence as shown in SEQ. ID. NO. 9;

8 v. a protein having an amino acid sequence as shown in SEQ. ID. NO. 11;
9 and

10 vi. any portion of any of the proteins above that elicits an immune response
11 against *E. canis*;

12 b) testing said peptide fragment to determine if said peptide fragment elicits an
13 immune response in a host animal; and

14 c) identifying said peptide fragment as said T cell epitope of *E. canis* if said
15 fragment elicits an immune response.

1 25. The method of claim 24 wherein said peptide fragment comprises nine to twenty
2 amino acids.

1 26. A method of creating a vaccine against *E. canis* comprising:

2 a) selecting a vector capable of expressing a recombinant DNA inserted into said
3 vector; and

4 b) inserting a recombinant DNA into said vector such that a recombinant protein
5 is expressed when said vector is provided in an appropriate host wherein said
6 DNA is selected from the group consisting of:

7 i. a recombinant DNA that encodes a protein having an amino acid
8 sequence as shown in SEQ. ID. NO. 3;

9 ii. a recombinant DNA that encodes a protein having an amino acid
10 sequence as shown in SEQ. ID. NO. 5;

11 iii. a recombinant DNA that encodes a protein having an amino acid
12 sequence as shown in SEQ. ID. NO. 7;

13 iv. a recombinant DNA that encodes a protein having an amino acid
14 sequence as shown in SEQ. ID. NO. 9;

15 v. a recombinant DNA that encodes a protein having an amino acid
16 sequence as shown in SEQ. ID. NO. 11; and

17 vi. any portion of said DNA above that encodes a protein that elicits an
18 immune response against *E. canis*.

1 27. The method of claim 26, wherein said DNA further comprises DNA that encodes CpG
2 motifs.

1 28. The method of claim 26 wherein said DNA further comprises a promoter selected from
2 the group consisting of:

3 a) a cytomegalovirus (CMV) immediate early promoter;

4 b) a human tissue plasminogen activator gene (t-PA); and

5 c) a promoter/enhancer region of a human elongation factor alpha (EF-1 α).

1 29. The method of claim 26, wherein said vector is selected from the group consisting of:

2 a) pcDNA3;

3 b) pC1;

4 c) VR1012; and

5 d) VR1020.

1 30. The method of claim 26 wherein said vaccine is injected into said host in a manner
2 selected from the group consisting of:

3 a) intramuscular injection;

4 b) intravenous injection; and

5 c) gene gun injection.

1 31. The method of claim 30, wherein said host is a dog.

1 32. A method of creating a vaccine against *E. canis* comprising:

2 a) selecting a vector capable of expressing a recombinant protein inserted into
3 said vector;

4 b) insertion of a recombinant DNA into said vector such that said recombinant
5 protein is expressed when said vector is transformed into a bacterial strain
6 wherein said DNA is selected from the group consisting of:

7 i. a recombinant DNA that encodes a protein having an amino acid
8 sequence as shown in SEQ. ID. NO. 3;

9 ii. a recombinant DNA that encodes a protein having an amino acid
10 sequence as shown in SEQ. ID. NO. 5;

- 11 iii. a recombinant DNA that encodes a protein having an amino acid
12 sequence as shown in SEQ. ID. NO. 7;
- 13 iv. a recombinant DNA that encodes a protein having an amino acid
14 sequence as shown in SEQ. ID. NO. 9;
- 15 v. a recombinant DNA that encodes a protein having an amino acid
16 sequence as shown in SEQ. ID. NO. 11; and
- 17 vi. any portion of said DNA above that encodes a protein that elicits an
18 immune response against *E. canis*; and
- 19 c) harvesting said recombinant protein from said bacterial strain.

1 33. The method of claim 32, wherein said vaccine further comprises adjuvants selected
2 from the group consisting of:

- 3 a) aluminum hydroxide;
- 4 b) QuilA; and
- 5 c) Montamide.

1 34. The method of claim 32, wherein said vaccine further comprises a promoter selected
2 from the group consisting of:

- 3 a) tac;
- 4 b) T5; and
- 5 c) T7.

1 35. The method of claim 32, wherein said bacterial strain is *E. coli*.

1 36. The method of claim 32, wherein said vector is selected from the group consisting of:

- 2 a) pREST;

3 b) pET; and

4 c) pKK233-3.

1 37. The method of claim 32 wherein said vaccine further comprises a cytokine operatively
2 associated with said vaccine.

1 38. The method of claim 37 wherein said cytokine is selected from the group consisting
2 of:

3 a) interleukin-1 β (IL-1 β);

4 b) granulocyte-macrophage colony stimulating factor (GM-CSF);

5 c) gamma interferon (γ -IFN);

6 d) amino acids VQGEESNDK from the IL-1 β protein; and

7 e) any portion of any of the cytokines above that elicits an improved
8 immunogenic response against *E. canis*.

1 39. The method of claim 32 wherein said vaccine is injected into said host in a manner
2 selected from the group consisting of:

3 a) intramuscular injection; and

4 b) subcutaneous injection.

1 40. The method of claim 39 wherein said host is a dog.

1 41. A method of creating a T cell epitope vaccine comprising:

2 a) selecting a recombinant protein that includes a T cell epitope wherein said T
3 cell epitope comprises an amino acid peptide fragment of a protein selected
4 from the group consisting of:

5 i. a protein having an amino acid sequence as shown in SEQ. ID. NO. 3;

- ii. a protein having an amino acid sequence as shown in SEQ. ID. NO. 5;
 - iii. a protein having an amino acid sequence as shown in SEQ. ID. NO. 7;
 - iv. a protein having an amino acid sequence as shown in SEQ. ID. NO. 9;
 - v. a protein having an amino acid sequence as shown in SEQ. ID. NO. 11;
and
 - vi. any portion of any of the above proteins that elicits an immune response
against *E. canis*;
- b) identifying said T cell epitope from said protein;
 - c) incorporating said T cell epitope into a construct capable of expressing said
epitope as a protein; and
 - d) harvesting said protein.

42. The method of claim 41 wherein said amino acid peptide fragment comprises nine to
twenty amino acids.

43. The method of claim 41 wherein said construct capable of expressing said epitope
further comprises a recombinant DNA encoding a protein which is capable of being
internalized into eukaryotic cells, including cells of the immune system.

44. The method of claim 43 wherein said protein capable of being internalized into
eukaryotic cells comprises a toxin selected from the group consisting of:

- a) a recombinant adenylate cyclase of *Bordetella bronchiseptica*; and
- b) a recombinant exotoxin A (PE) of *Pseudomonas aeruginosa*.

45. The method of claim 41 wherein said vaccine is injected into said host in a manner
selected from the group consisting of:


- a) intramuscular injection; and

4 b) subcutaneous injection.

1 46. The method of claim 45 wherein said host is a dog.

1/1

Figure 1



SEQUENCE LISTING

<110> Chang, Yung-Fu

<120> EHRlichia CANIS GENES FOR VACCINE DEVELOPMENT

<130> crf2322

<150> U.S. 09/358,322

<151> 1999-07-21

<160> 11

<170> PatentIn Ver. 2.0

<210> 1

<211> 5300

<212> DNA

<213> Ehrlichia canis

<400> 1

gatcaaataa aatgaaacca agaataagaa acactatttta tggattaata gcaataatac 60
tatctatgat atgttttagtg tacgcttctg taccactata tagtatattt tgtaaagtaa 120
cagggttatgg aggtacagta agaacaagta atatatcaaa ttctaaaata ggtaacacta 180
ttattaaagt cagatttaaat gcagatatac acaaacaact gccatggaaa ttctatccag 240
aagtatctca tgtatttgta aaaccaggag aacaaaaatt gatcttctac cgcgagaaaa 300
atctacttga tgaggacact tcaggaatgg ctgtatataa tgttacacca cataaagtag 360
gaaaatattt taataaggta gcttggtttt gtttcaccaa acaaacatta taccctcatc 420
aaaaaactat aatgccagta tcatttttta tagatccagc catagaaaca gatcctgaaa 480
ctgctgacgt aaaactcatc actctttcat atgtattctt taagtacaaa gaataaactt 540
catataccgt acattataaa ctgattaaaa aaaataacta ttaatatga gcaaaataat 600
ttatctattc aacagattct tttcaattag agagtattca aaaacactac aactactgct 660
tgcaactttc tatcactgat atataaaagt gaaataaatt taaaaaactt tagttttaat 720
agaagaattt tattaaaagc tttgaatcaa atttaattac tgatataaaa atactattaa 780
acattaacaa tgcttaatta aagtattatt atttacctta atttcataac ctttattaac 840

aatttcataa taaaaataact ttactctttat ttttttatca cttgatatta ttaaataatc 900
atataaactc ccaaataaac tattgcaagg ttatggtaat gatgaaattt tttacttggt 960
ttttcatagt tttcttaaca atagccaatc atgctttatc ctttaacatt aaagttacac 1020
atgaaaaatt agataatgga atggaagtat acgtgattcc aaatcatcgc gcaccagcag 1080
tcatgcacat ggtattatac aaagtcggtg gaactgatga tccagtagga tactctggat 1140
tagcacattt ttttgaacac ttaatgttta gtggaacaga aaaatttcct aatctcatca 1200
gcacacttag taatataggc ggaaatttca atgcaagcac atctcaattt tgtactatat 1260
actacgaatt aataccaaaa caatatttat ctcttgcaat ggatattgaa tcagacagaa 1320
tgcagaattt taaggttacc gacaaagcat taataagaga acaaaaggta gtcttagaag 1380
aaagaaaaat gagagttgaa agccaagcaa aaaacatact agaagaagaa atggaaaatg 1440
cattttatta caatggatat ggcagaccag tagtaggatg ggaacatgaa attagcaact 1500
acaacaaaga agttgctgaa gcctttcata agctacatta tagtcctaata aatgctatat 1560
taattgtaac tggagatgca gatccacaag aagtaatcac acttgcaaaa caatactatg 1620
ggaaaatacc atctaataat aagaaacctt caagtcaagt tagggtagaa ccaccgcata 1680
aaacaaatat gactttaaca ttaaaagaca gttcagtaga aatcccagaa ctgtttttta 1740
tgtatcaaata accaaatggg attaccaata aaaactacat acttaacatg atgttagcag 1800
aaatactcgg tagtggtaaa ttcagcctgc tttacaatga tttggtaatt aacaatccaa 1860
tagttacatc gataaaaaca gattataatt acttaactga cagcgataat tacctttcca 1920
ttgaagctat acctaaaaac gggatctcta cagaagctgt agaacaagaa attcataaat 1980
gtataaataa ttatttagaa aatggaattt cagcagaata tttagaaagt gcaaagtata 2040
aagtaaaagc acattttaact tatgcatttg acggactaac tttcatatca tatttttatg 2100
gcatgcatct aatactagga gtaccgctat cagaaatcag taatatttac gataccatag 2160
acaaagtaag tatccaagat gttaactccg ctatggaaaa tatctttcaa aacaatataa 2220
gattaaccgg gcatttatta cctaattggag aatagttatg agaaacatat tgtgttacac 2280
attaatattg attttctttt cattcaatac atatgcaaat gatctcaata ttaacataaa 2340
agaagctaca actaaaaata aaatacacta tctatatgtt gaacatcata acctaccaac 2400
aatttcctta aaatttgcac tcaagaaagc aggatacgct tatgatgcct ttgataagca 2460
aggacttgca tactttacat caaaaatatt aaacgaagga tcaaaaaaca actatgctct 2520
cagttttgca caacaattag aaggcaaagg tatagactta aaatttgata tagacctaga 2580

caatTTTTat atatcattaa aaaccttatc agaaaacttt gaagaagccc tagttttact 2640
cagtgattgc atattcaaca ccgtcacaga tcaagaaata ttcaatagaa taatagcaga 2700
acagattgca catgttaa at cattatatc tgctcctgaa tttatagcta caacagaaat 2760
gaatcacgct atattcaaag ggcaccata ttctaacaaa gtttacggga cattaaatac 2820
aatcaataat atcaaccagg aagacgttgc attatatata aaaaatagtt ttgacaagga 2880
acaaatcggt atcagcgcag caggagatgt agatccaaca cagctatcaa atttactaga 2940
taaataatatt ctttccaaat tgccatctgg taataacaaa aataccatac cagatacgac 3000
tgtaataaga gaagacacat tattatatgt acagagagat gtaccacaaa gtgtcataat 3060
gtttgctaca gacacagtac catatcacag caaagactat catgcatcaa acttggtcaa 3120
tactatgcta ggcggattaa gtctcaattc aatattaatg atagaattaa gagacaagtt 3180
aggattaaca taccatagta gcagttcact atctaactg aatcatagta atgtgctatt 3240
tggtacaata ttactgata ataccacagt aacaaaatgt atatccgtct taacagatat 3300
tatagagcac attaaaaagt atggagttga tgaagacact tttgcaattg caaaatctag 3360
tattaccaac tcttttattt tatctatgtt aaataacaat aatgttagtg agatattgtt 3420
aagcttacaa ttacacgac tagatccgag ttatattaat aaatacaatt cttactacaa 3480
agcaataaca atagaagaag taaataaaat tgccaagaaa attttatcta atgaattagt 3540
aataattgaa gtaggaaaaa acaataacat aaatggcaaa caaatagatg ctaaaaaaca 3600
cataccttgg ttaagtatac aggttattgt atttactaca agtattctat taggttgat 3660
taagtaagta taagtagctt caatcaaata aaaaaacatt aaccaaagtg ttagctctac 3720
cggagaagct tattataagc ttttaacctg ggataatatg aagttttgct aatgttaagc 3780
aaaaaattag taatcacaat atcaaatttt ctttacagga ttatattgtg acctaccata 3840
acaacttata tttagaaaat gacaacagat acacacatca ataaattatc actacaattc 3900
aattaataaaa acaatgagta tttttactta attatttaat tttatttttt aaaataaaat 3960
tacaatttta cttactcaat aaaagcagtt atactaccaa gtattggatg gtattaatcg 4020
gagcaattac tacttaatag tatagctgtt gacaagccgc aatctgcggt tcttgacaaa 4080
ataatactaa tcagttaaaa ttttgaagtg tttcaccata atgggtattat ttatgaaagc 4140
tcatagcaca agtatacgga actttcagcc tttagaaaga gctgctataa tcattgcagt 4200
gttaggttta gctgcattct tgtttgctgc tgctgcctgc agtgatcggt tccaaagatt 4260

gcaattaaca aatccatttg taatagcagg aatgggttggc cttgcagttc ttttagttgc 4320
ttccttaaca gcagcattaa gtatatgctt aactaaaagt aagcaagtca cacaacatgc 4380
tattagacat cgcttttgat acgagtcaag cacttcttct tctgtactgc ttgcaatatc 4440
aataatttct ttattacttg ctgcagcatt ttgtggaaag ataatgggta atgacaaccc 4500
agatctattc ttttagcaaga tgcaagaact ctccaatcca cttgttggtg cagctattgt 4560
agccgtttct gttttcctac tctcattcgt aatgtatgct gcaaagaaca ttataagtcc 4620
agataaacia actcacgtta ttatattatc taatcaacia actatagaag aagcaaaagt 4680
agatcaagga atgaatattt tgtcagcagt actcccagca gctggcattg acatcatgac 4740
tatagcttct tgtgacattt tagcagttag cagccgggga tcctctcagc atcaatagat 4800
ttatgtttta gcctgtattc acctttttat taggtgttgt atcgtttctt tatataagtg 4860
tggtatatta tataaaacat ctaggagtta cagttaattt gtttcatgtg gttattactc 4920
tttgccatta ttattactat acctaaaaat ataaaagaat ccgccagggt gaatacaggc 4980
caatgtaagt tattgatata aaaatctata aaatcataga cagcaccata tcttattcta 5040
tctatgatat ttcctattga ccccccaata atgattacaa gaggtaatct ataatgtggc 5100
tgtactataa ataagtagca taaaacacia gtaatcaaaa tcgagatact acaaaaaaca 5160
acattactat attcaaagtt atttaataata caaaactaa ttccagcatt ccacactgta 5220
gtaaagcgca agaagcttaa tatctctatt acacctttat ctcctatcaa atttactaca 5280
taccatttac ttacctgatc 5300

<210> 2

<211> 522

<212> DNA

<213> Ehrlichia canis

<220>

<221> CDS

<222> (1)..(522)

<223> Protein translated from nucleotides 12 through 533
(cytochrome oxidase homolog).

<400> 2

5

atg	aaa	cca	aga	ata	aga	aac	act	att	tat	gga	tta	ata	gca	ata	ata	48
Met	Lys	Pro	Arg	Ile	Arg	Asn	Thr	Ile	Tyr	Gly	Leu	Ile	Ala	Ile	Ile	
1				5				10					15			
cta	tct	atg	ata	tgt	tta	gtg	tac	gct	tct	gta	cca	cta	tat	agt	ata	96
Leu	Ser	Met	Ile	Cys	Leu	Val	Tyr	Ala	Ser	Val	Pro	Leu	Tyr	Ser	Ile	
			20					25					30			
ttt	tgt	aaa	gta	aca	ggg	tat	gga	ggg	aca	gta	aga	aca	agt	aat	ata	144
Phe	Cys	Lys	Val	Thr	Gly	Tyr	Gly	Gly	Thr	Val	Arg	Thr	Ser	Asn	Ile	
		35					40					45				
tca	aat	tct	aaa	ata	ggg	aac	act	att	att	aaa	gtc	aga	ttt	aat	gca	192
Ser	Asn	Ser	Lys	Ile	Gly	Asn	Thr	Ile	Ile	Lys	Val	Arg	Phe	Asn	Ala	
	50					55					60					
gat	ata	cac	aaa	caa	ctg	cca	tgg	aaa	ttc	tat	cca	gaa	gta	tct	cat	240
Asp	Ile	His	Lys	Gln	Leu	Pro	Trp	Lys	Phe	Tyr	Pro	Glu	Val	Ser	His	
	65				70					75					80	
gta	ttt	gta	aaa	cca	gga	gaa	caa	aaa	ttg	att	ttc	tac	cgc	gca	gaa	288
Val	Phe	Val	Lys	Pro	Gly	Glu	Gln	Lys	Leu	Ile	Phe	Tyr	Arg	Ala	Glu	
			85					90						95		
aat	cta	ctt	gat	gag	gac	act	tca	gga	atg	gct	gta	tat	aat	gtt	aca	336
Asn	Leu	Leu	Asp	Glu	Asp	Thr	Ser	Gly	Met	Ala	Val	Tyr	Asn	Val	Thr	
			100					105					110			
cca	cat	aaa	gta	gga	aaa	tat	ttt	aat	aag	gta	gct	tgt	ttt	tgt	ttc	384
Pro	His	Lys	Val	Gly	Lys	Tyr	Phe	Asn	Lys	Val	Ala	Cys	Phe	Cys	Phe	
		115					120					125				
acc	aaa	caa	aca	tta	tac	cct	cat	caa	aaa	act	ata	atg	cca	gta	tca	432
Thr	Lys	Gln	Thr	Leu	Tyr	Pro	His	Gln	Lys	Thr	Ile	Met	Pro	Val	Ser	
		130				135					140					
ttt	ttt	ata	gat	cca	gcc	ata	gaa	aca	gat	cct	gaa	act	gct	gac	gta	480
Phe	Phe	Ile	Asp	Pro	Ala	Ile	Glu	Thr	Asp	Pro	Glu	Thr	Ala	Asp	Val	
145					150					155					160	
aaa	ctc	atc	act	ctt	tca	tat	gta	ttc	ttt	aag	tac	aaa	gaa			522
Lys	Leu	Ile	Thr	Leu	Ser	Tyr	Val	Phe	Phe	Lys	Tyr	Lys	Glu			
				165					170							

<210> 3

<211> 174

<212> PRT

<213> Ehrlichia canis

<400> 3

Met	Lys	Pro	Arg	Ile	Arg	Asn	Thr	Ile	Tyr	Gly	Leu	Ile	Ala	Ile	Ile
1				5					10					15	

Leu Ser Met Ile Cys Leu Val Tyr Ala Ser Val Pro Leu Tyr Ser Ile
 20 25 30
 Phe Cys Lys Val Thr Gly Tyr Gly Gly Thr Val Arg Thr Ser Asn Ile
 35 40 45
 Ser Asn Ser Lys Ile Gly Asn Thr Ile Ile Lys Val Arg Phe Asn Ala
 50 55 60
 Asp Ile His Lys Gln Leu Pro Trp Lys Phe Tyr Pro Glu Val Ser His
 65 70 75 80
 Val Phe Val Lys Pro Gly Glu Gln Lys Leu Ile Phe Tyr Arg Ala Glu
 85 90 95
 Asn Leu Leu Asp Glu Asp Thr Ser Gly Met Ala Val Tyr Asn Val Thr
 100 105 110
 Pro His Lys Val Gly Lys Tyr Phe Asn Lys Val Ala Cys Phe Cys Phe
 115 120 125
 Thr Lys Gln Thr Leu Tyr Pro His Gln Lys Thr Ile Met Pro Val Ser
 130 135 140
 Phe Phe Ile Asp Pro Ala Ile Glu Thr Asp Pro Glu Thr Ala Asp Val
 145 150 155 160
 Lys Leu Ile Thr Leu Ser Tyr Val Phe Phe Lys Tyr Lys Glu
 165 170

<210> 4

<211> 1314

<212> DNA

<213> Ehrlichia canis

<220>

<221> CDS

<222> (1)..(1314)

<223> Protein translated from nucleotides 939 through
 2,252 (ProA).

<400> 4

atg atg aaa ttt ttt act tgt ttt ttc ata gtt ttc tta aca ata gcc 48
 Met Met Lys Phe Phe Thr Cys Phe Phe Ile Val Phe Leu Thr Ile Ala
 1 5 10 15

7

aat cat gct tta tcc ttt aac att aaa gtt aca cat gaa aaa tta gat	96
Asn His Ala Leu Ser Phe Asn Ile Lys Val Thr His Glu Lys Leu Asp	
20 25 30	
aat gga atg gaa gta tac gtg att cca aat cat cgc gca cca gca gtc	144
Asn Gly Met Glu Val Tyr Val Ile Pro Asn His Arg Ala Pro Ala Val	
35 40 45	
atg cac atg gta tta tac aaa gtc ggt gga act gat gat cca gta gga	192
Met His Met Val Leu Tyr Lys Val Gly Gly Thr Asp Asp Pro Val Gly	
50 55 60	
tac tct gga tta gca cat ttt ttt gaa cac tta atg ttt agt gga aca	240
Tyr Ser Gly Leu Ala His Phe Phe Glu His Leu Met Phe Ser Gly Thr	
65 70 75 80	
gaa aaa ttt cct aat ctc atc agc aca ctt agt aat ata ggc gga aat	288
Glu Lys Phe Pro Asn Leu Ile Ser Thr Leu Ser Asn Ile Gly Gly Asn	
85 90 95	
ttc aat gca agc aca tct caa ttt tgt act ata tac tac gaa tta ata	336
Phe Asn Ala Ser Thr Ser Gln Phe Cys Thr Ile Tyr Tyr Glu Leu Ile	
100 105 110	
cca aaa caa tat tta tct ctt gca atg gat att gaa tca gac aga atg	384
Pro Lys Gln Tyr Leu Ser Leu Ala Met Asp Ile Glu Ser Asp Arg Met	
115 120 125	
cag aat ttt aag gtt acc gac aaa gca tta ata aga gaa caa aag gta	432
Gln Asn Phe Lys Val Thr Asp Lys Ala Leu Ile Arg Glu Gln Lys Val	
130 135 140	
gtc tta gaa gaa aga aaa atg aga gtt gaa agc caa gca aaa aac ata	480
Val Leu Glu Glu Arg Lys Met Arg Val Glu Ser Gln Ala Lys Asn Ile	
145 150 155 160	
cta gaa gaa gaa atg gaa aat gca ttt tat tac aat gga tat ggc aga	528
Leu Glu Glu Glu Met Glu Asn Ala Phe Tyr Tyr Asn Gly Tyr Gly Arg	
165 170 175	
cca gta gta gga tgg gaa cat gaa att agc aac tac aac aaa gaa gtt	576
Pro Val Val Gly Trp Glu His Glu Ile Ser Asn Tyr Asn Lys Glu Val	
180 185 190	
gct gaa gcc ttt cat aag cta cat tat agt cct aat aat gct ata tta	624
Ala Glu Ala Phe His Lys Leu His Tyr Ser Pro Asn Asn Ala Ile Leu	
195 200 205	
att gta act gga gat gca gat cca caa gaa gta atc aca ctt gca aaa	672
Ile Val Thr Gly Asp Ala Asp Pro Gln Glu Val Ile Thr Leu Ala Lys	
210 215 220	
caa tac tat ggg aaa ata cca tct aat aat aag aaa cct tca agt caa	720
Gln Tyr Tyr Gly Lys Ile Pro Ser Asn Asn Lys Lys Pro Ser Ser Gln	
225 230 235 240	

gtt agg gta gaa cca ccg cat aaa aca aat atg act tta aca tta aaa	768
Val Arg Val Glu Pro Pro His Lys Thr Asn Met Thr Leu Thr Leu Lys	
245 250 255	
gac agt tca gta gaa atc cca gaa ctg ttt tta atg tat caa ata cca	816
Asp Ser Ser Val Glu Ile Pro Glu Leu Phe Leu Met Tyr Gln Ile Pro	
260 265 270	
aat ggt att acc aat aaa aac tac ata ctt aac atg atg tta gca gaa	864
Asn Gly Ile Thr Asn Lys Asn Tyr Ile Leu Asn Met Met Leu Ala Glu	
275 280 285	
ata ctc ggt agt ggt aaa ttc agc ctg ctt tac aat gat ttg gta att	912
Ile Leu Gly Ser Gly Lys Phe Ser Leu Leu Tyr Asn Asp Leu Val Ile	
290 295 300	
aac aat cca ata gtt aca tcg ata aaa aca gat tat aat tac tta act	960
Asn Asn Pro Ile Val Thr Ser Ile Lys Thr Asp Tyr Asn Tyr Leu Thr	
305 310 315 320	
gac agc gat aat tac ctt tcc att gaa gct ata cct aaa aac ggg atc	1008
Asp Ser Asp Asn Tyr Leu Ser Ile Glu Ala Ile Pro Lys Asn Gly Ile	
325 330 335	
tct aca gaa gct gta gaa caa gaa att cat aaa tgt ata aat aat tat	1056
Ser Thr Glu Ala Val Glu Gln Glu Ile His Lys Cys Ile Asn Asn Tyr	
340 345 350	
tta gaa aat gga att tca gca gaa tat tta gaa agt gca aag tat aaa	1104
Leu Glu Asn Gly Ile Ser Ala Glu Tyr Leu Glu Ser Ala Lys Tyr Lys	
355 360 365	
gta aaa gca cat tta act tat gca ttt gac gga cta act ttc ata tca	1152
Val Lys Ala His Leu Thr Tyr Ala Phe Asp Gly Leu Thr Phe Ile Ser	
370 375 380	
tat ttt tat ggc atg cat cta ata cta gga gta ccg cta tca gaa atc	1200
Tyr Phe Tyr Gly Met His Leu Ile Leu Gly Val Pro Leu Ser Glu Ile	
385 390 395 400	
agt aat att tac gat acc ata gac aaa gta agt atc caa gat gtt aac	1248
Ser Asn Ile Tyr Asp Thr Ile Asp Lys Val Ser Ile Gln Asp Val Asn	
405 410 415	
tcc gct atg gaa aat atc ttt caa aac aat ata aga tta acc ggg cat	1296
Ser Ala Met Glu Asn Ile Phe Gln Asn Asn Ile Arg Leu Thr Gly His	
420 425 430	
tta tta cct aat gga gaa	1314
Leu Leu Pro Asn Gly Glu	
435	

<210> 5

<211> 438

<212> PRT

<213> Ehrlichia canis

<400> 5

Met	Met	Lys	Phe	Phe	Thr	Cys	Phe	Phe	Ile	Val	Phe	Leu	Thr	Ile	Ala	1	5	10	15
Asn	His	Ala	Leu	Ser	Phe	Asn	Ile	Lys	Val	Thr	His	Glu	Lys	Leu	Asp	20	25	30	
Asn	Gly	Met	Glu	Val	Tyr	Val	Ile	Pro	Asn	His	Arg	Ala	Pro	Ala	Val	35	40	45	
Met	His	Met	Val	Leu	Tyr	Lys	Val	Gly	Gly	Thr	Asp	Asp	Pro	Val	Gly	50	55	60	
Tyr	Ser	Gly	Leu	Ala	His	Phe	Phe	Glu	His	Leu	Met	Phe	Ser	Gly	Thr	65	70	75	80
Glu	Lys	Phe	Pro	Asn	Leu	Ile	Ser	Thr	Leu	Ser	Asn	Ile	Gly	Gly	Asn	85	90	95	
Phe	Asn	Ala	Ser	Thr	Ser	Gln	Phe	Cys	Thr	Ile	Tyr	Tyr	Glu	Leu	Ile	100	105	110	
Pro	Lys	Gln	Tyr	Leu	Ser	Leu	Ala	Met	Asp	Ile	Glu	Ser	Asp	Arg	Met	115	120	125	
Gln	Asn	Phe	Lys	Val	Thr	Asp	Lys	Ala	Leu	Ile	Arg	Glu	Gln	Lys	Val	130	135	140	
Val	Leu	Glu	Glu	Arg	Lys	Met	Arg	Val	Glu	Ser	Gln	Ala	Lys	Asn	Ile	145	150	155	160
Leu	Glu	Glu	Glu	Met	Glu	Asn	Ala	Phe	Tyr	Tyr	Asn	Gly	Tyr	Gly	Arg	165	170	175	
Pro	Val	Val	Gly	Trp	Glu	His	Glu	Ile	Ser	Asn	Tyr	Asn	Lys	Glu	Val	180	185	190	
Ala	Glu	Ala	Phe	His	Lys	Leu	His	Tyr	Ser	Pro	Asn	Asn	Ala	Ile	Leu	195	200	205	
Ile	Val	Thr	Gly	Asp	Ala	Asp	Pro	Gln	Glu	Val	Ile	Thr	Leu	Ala	Lys	210	215	220	
Gln	Tyr	Tyr	Gly	Lys	Ile	Pro	Ser	Asn	Asn	Lys	Lys	Pro	Ser	Ser	Gln	225	230	235	240
Val	Arg	Val	Glu	Pro	Pro	His	Lys	Thr	Asn	Met	Thr	Leu	Thr	Leu	Lys	245	250	255	
Asp	Ser	Ser	Val	Glu	Ile	Pro	Glu	Leu	Phe	Leu	Met	Tyr	Gln	Ile	Pro	260	265	270	

10

Asn Gly Ile Thr Asn Lys Asn Tyr Ile Leu Asn Met Met Leu Ala Glu
 275 280 285
 Ile Leu Gly Ser Gly Lys Phe Ser Leu Leu Tyr Asn Asp Leu Val Ile
 290 295 300
 Asn Asn Pro Ile Val Thr Ser Ile Lys Thr Asp Tyr Asn Tyr Leu Thr
 305 310 315 320
 Asp Ser Asp Asn Tyr Leu Ser Ile Glu Ala Ile Pro Lys Asn Gly Ile
 325 330 335
 Ser Thr Glu Ala Val Glu Gln Glu Ile His Lys Cys Ile Asn Asn Tyr
 340 345 350
 Leu Glu Asn Gly Ile Ser Ala Glu Tyr Leu Glu Ser Ala Lys Tyr Lys
 355 360 365
 Val Lys Ala His Leu Thr Tyr Ala Phe Asp Gly Leu Thr Phe Ile Ser
 370 375 380
 Tyr Phe Tyr Gly Met His Leu Ile Leu Gly Val Pro Leu Ser Glu Ile
 385 390 395 400
 Ser Asn Ile Tyr Asp Thr Ile Asp Lys Val Ser Ile Gln Asp Val Asn
 405 410 415
 Ser Ala Met Glu Asn Ile Phe Gln Asn Asn Ile Arg Leu Thr Gly His
 420 425 430
 Leu Leu Pro Asn Gly Glu
 435

<210> 6

<211> 1407

<212> DNA

<213> Ehrlichia canis

<220>

<221> CDS

<222> (1)..(1407)

<223> Protein translated from 2,258 through 3,664
 (ProB) .

<400> 6

atg aga aac ata ttg tgt tac aca tta ata ttg att ttc ttt tca ttc 48
 Met Arg Asn Ile Leu Cys Tyr Thr Leu Ile Leu Ile Phe Phe Ser Phe
 1 5 10 15

aat	aca	tat	gca	aat	gat	ctc	aat	att	aac	ata	aaa	gaa	gct	aca	act	96
Asn	Thr	Tyr	Ala	Asn	Asp	Leu	Asn	Ile	Asn	Ile	Lys	Glu	Ala	Thr	Thr	
			20					25					30			
aaa	aat	aaa	ata	cac	tat	cta	tat	gtt	gaa	cat	cat	aac	cta	cca	aca	144
Lys	Asn	Lys	Ile	His	Tyr	Leu	Tyr	Val	Glu	His	His	Asn	Leu	Pro	Thr	
		35					40					45				
att	tcc	tta	aaa	ttt	gca	ttc	aag	aaa	gca	gga	tac	gct	tat	gat	gcc	192
Ile	Ser	Leu	Lys	Phe	Ala	Phe	Lys	Lys	Ala	Gly	Tyr	Ala	Tyr	Asp	Ala	
	50					55					60					
ttt	gat	aag	caa	gga	ctt	gca	tac	ttt	aca	tca	aaa	ata	tta	aac	gaa	240
Phe	Asp	Lys	Gln	Gly	Leu	Ala	Tyr	Phe	Thr	Ser	Lys	Ile	Leu	Asn	Glu	
65					70					75					80	
gga	tca	aaa	aac	aac	tat	gct	ctc	agt	ttt	gca	caa	caa	tta	gaa	ggc	288
Gly	Ser	Lys	Asn	Asn	Tyr	Ala	Leu	Ser	Phe	Ala	Gln	Gln	Leu	Glu	Gly	
			85					90						95		
aaa	ggt	ata	gac	tta	aaa	ttt	gat	ata	gac	cta	gac	aat	ttt	tat	ata	336
Lys	Gly	Ile	Asp	Leu	Lys	Phe	Asp	Ile	Asp	Leu	Asp	Asn	Phe	Tyr	Ile	
			100					105					110			
tca	tta	aaa	acc	tta	tca	gaa	aac	ttt	gaa	gaa	gcc	cta	gtt	tta	ctc	384
Ser	Leu	Lys	Thr	Leu	Ser	Glu	Asn	Phe	Glu	Glu	Ala	Leu	Val	Leu	Leu	
		115					120					125				
agt	gat	tgc	ata	ttc	aac	acc	gtc	aca	gat	caa	gaa	ata	ttc	aat	aga	432
Ser	Asp	Cys	Ile	Phe	Asn	Thr	Val	Thr	Asp	Gln	Glu	Ile	Phe	Asn	Arg	
	130					135					140					
ata	ata	gca	gaa	cag	att	gca	cat	gtt	aaa	tca	tta	tat	tct	gct	cct	480
Ile	Ile	Ala	Glu	Gln	Ile	Ala	His	Val	Lys	Ser	Leu	Tyr	Ser	Ala	Pro	
145					150					155					160	
gaa	ttt	ata	gct	aca	aca	gaa	atg	aat	cac	gct	ata	ttc	aaa	ggg	cac	528
Glu	Phe	Ile	Ala	Thr	Thr	Glu	Met	Asn	His	Ala	Ile	Phe	Lys	Gly	His	
				165					170					175		
cca	tat	tct	aac	aaa	gtt	tac	ggg	aca	tta	aat	aca	atc	aat	aat	atc	576
Pro	Tyr	Ser	Asn	Lys	Val	Tyr	Gly	Thr	Leu	Asn	Thr	Ile	Asn	Asn	Ile	
			180					185					190			
aac	cag	gaa	gac	gtt	gca	tta	tat	ata	aaa	aat	agt	ttt	gac	aag	gaa	624
Asn	Gln	Glu	Asp	Val	Ala	Leu	Tyr	Ile	Lys	Asn	Ser	Phe	Asp	Lys	Glu	
		195					200					205				
caa	atc	gtt	atc	agc	gca	gca	gga	gat	gta	gat	cca	aca	cag	cta	tca	672
Gln	Ile	Val	Ile	Ser	Ala	Ala	Gly	Asp	Val	Asp	Pro	Thr	Gln	Leu	Ser	
	210					215					220					
aat	tta	cta	gat	aaa	tat	att	ctt	tcc	aaa	ttg	cca	tct	ggg	aat	aac	720
Asn	Leu	Leu	Asp	Lys	Tyr	Ile	Leu	Ser	Lys	Leu	Pro	Ser	Gly	Asn	Asn	
225					230					235					240	

12

aaa aat acc ata cca gat acg act gtt aat aga gaa gac aca tta tta	768
Lys Asn Thr Ile Pro Asp Thr Thr Val Asn Arg Glu Asp Thr Leu Leu	
245 250 255	
tat gta cag aga gat gta cca caa agt gtc ata atg ttt gct aca gac	816
Tyr Val Gln Arg Asp Val Pro Gln Ser Val Ile Met Phe Ala Thr Asp	
260 265 270	
aca gta cca tat cac agc aaa gac tat cat gca tca aac ttg ttc aat	864
Thr Val Pro Tyr His Ser Lys Asp Tyr His Ala Ser Asn Leu Phe Asn	
275 280 285	
act atg cta ggc gga tta agt ctc aat tca ata tta atg ata gaa tta	912
Thr Met Leu Gly Gly Leu Ser Leu Asn Ser Ile Leu Met Ile Glu Leu	
290 295 300	
aga gac aag tta gga tta aca tac cat agt agc agt tca cta tct aac	960
Arg Asp Lys Leu Gly Leu Thr Tyr His Ser Ser Ser Ser Leu Ser Asn	
305 310 315 320	
atg aat cat agt aat gtg cta ttt ggt aca ata ttc act gat aat acc	1008
Met Asn His Ser Asn Val Leu Phe Gly Thr Ile Phe Thr Asp Asn Thr	
325 330 335	
aca gta aca aaa tgt ata tcc gtc tta aca gat att ata gag cac att	1056
Thr Val Thr Lys Cys Ile Ser Val Leu Thr Asp Ile Ile Glu His Ile	
340 345 350	
aaa aag tat gga gtt gat gaa gac act ttt gca att gca aaa tct agt	1104
Lys Lys Tyr Gly Val Asp Glu Asp Thr Phe Ala Ile Ala Lys Ser Ser	
355 360 365	
att acc aac tct ttt att tta tct atg tta aat aac aat aat gtt agt	1152
Ile Thr Asn Ser Phe Ile Leu Ser Met Leu Asn Asn Asn Asn Val Ser	
370 375 380	
gag ata ttg tta agc tta caa tta cac gat cta gat ccg agt tat att	1200
Glu Ile Leu Leu Ser Leu Gln Leu His Asp Leu Asp Pro Ser Tyr Ile	
385 390 395 400	
aat aaa tac aat tct tac tac aaa gca ata aca ata gaa gaa gta aat	1248
Asn Lys Tyr Asn Ser Tyr Tyr Lys Ala Ile Thr Ile Glu Glu Val Asn	
405 410 415	
aaa att gcc aag aaa att tta tct aat gaa tta gta ata att gaa gta	1296
Lys Ile Ala Lys Lys Ile Leu Ser Asn Glu Leu Val Ile Ile Glu Val	
420 425 430	
gga aaa aac aat aac ata aat ggc aaa caa ata gat gct aaa aaa cac	1344
Gly Lys Asn Asn Asn Ile Asn Gly Lys Gln Ile Asp Ala Lys Lys His	
435 440 445	
ata cct tgg tta agt ata cag gtt att gta ttt act aca agt att cta	1392
Ile Pro Trp Leu Ser Ile Gln Val Ile Val Phe Thr Thr Ser Ile Leu	
450 455 460	

tta ggt tgt att aag
 Leu Gly Cys Ile Lys
 465

1407

<210> 7

<211> 469

<212> PRT

<213> *Ehrlichia canis*

<400> 7

Met	Arg	Asn	Ile	Leu	Cys	Tyr	Thr	Leu	Ile	Leu	Ile	Phe	Phe	Ser	Phe		
1				5					10					15			
Asn	Thr	Tyr	Ala	Asn	Asp	Leu	Asn	Ile	Asn	Ile	Lys	Glu	Ala	Thr	Thr		
			20					25					30				
Lys	Asn	Lys	Ile	His	Tyr	Leu	Tyr	Val	Glu	His	His	Asn	Leu	Pro	Thr		
		35					40					45					
Ile	Ser	Leu	Lys	Phe	Ala	Phe	Lys	Lys	Ala	Gly	Tyr	Ala	Tyr	Asp	Ala		
	50					55					60						
Phe	Asp	Lys	Gln	Gly	Leu	Ala	Tyr	Phe	Thr	Ser	Lys	Ile	Leu	Asn	Glu		
65					70					75					80		
Gly	Ser	Lys	Asn	Asn	Tyr	Ala	Leu	Ser	Phe	Ala	Gln	Gln	Leu	Glu	Gly		
				85					90					95			
Lys	Gly	Ile	Asp	Leu	Lys	Phe	Asp	Ile	Asp	Leu	Asp	Asn	Phe	Tyr	Ile		
			100					105					110				
Ser	Leu	Lys	Thr	Leu	Ser	Glu	Asn	Phe	Glu	Glu	Ala	Leu	Val	Leu	Leu		
		115					120					125					
Ser	Asp	Cys	Ile	Phe	Asn	Thr	Val	Thr	Asp	Gln	Glu	Ile	Phe	Asn	Arg		
	130					135					140						
Ile	Ile	Ala	Glu	Gln	Ile	Ala	His	Val	Lys	Ser	Leu	Tyr	Ser	Ala	Pro		
145					150					155					160		
Glu	Phe	Ile	Ala	Thr	Thr	Glu	Met	Asn	His	Ala	Ile	Phe	Lys	Gly	His		
				165					170					175			
Pro	Tyr	Ser	Asn	Lys	Val	Tyr	Gly	Thr	Leu	Asn	Thr	Ile	Asn	Asn	Ile		
			180					185					190				
Asn	Gln	Glu	Asp	Val	Ala	Leu	Tyr	Ile	Lys	Asn	Ser	Phe	Asp	Lys	Glu		
		195					200					205					
Gln	Ile	Val	Ile	Ser	Ala	Ala	Gly	Asp	Val	Asp	Pro	Thr	Gln	Leu	Ser		
	210					215					220						

14

Asn Leu Leu Asp Lys Tyr Ile Leu Ser Lys Leu Pro Ser Gly Asn Asn
 225 230 235 240
 Lys Asn Thr Ile Pro Asp Thr Thr Val Asn Arg Glu Asp Thr Leu Leu
 245 250 255
 Tyr Val Gln Arg Asp Val Pro Gln Ser Val Ile Met Phe Ala Thr Asp
 260 265 270
 Thr Val Pro Tyr His Ser Lys Asp Tyr His Ala Ser Asn Leu Phe Asn
 275 280 285
 Thr Met Leu Gly Gly Leu Ser Leu Asn Ser Ile Leu Met Ile Glu Leu
 290 295 300
 Arg Asp Lys Leu Gly Leu Thr Tyr His Ser Ser Ser Ser Leu Ser Asn
 305 310 315 320
 Met Asn His Ser Asn Val Leu Phe Gly Thr Ile Phe Thr Asp Asn Thr
 325 330 335
 Thr Val Thr Lys Cys Ile Ser Val Leu Thr Asp Ile Ile Glu His Ile
 340 345 350
 Lys Lys Tyr Gly Val Asp Glu Asp Thr Phe Ala Ile Ala Lys Ser Ser
 355 360 365
 Ile Thr Asn Ser Phe Ile Leu Ser Met Leu Asn Asn Asn Asn Val Ser
 370 375 380
 Glu Ile Leu Leu Ser Leu Gln Leu His Asp Leu Asp Pro Ser Tyr Ile
 385 390 395 400
 Asn Lys Tyr Asn Ser Tyr Tyr Lys Ala Ile Thr Ile Glu Glu Val Asn
 405 410 415
 Lys Ile Ala Lys Lys Ile Leu Ser Asn Glu Leu Val Ile Ile Glu Val
 420 425 430
 Gly Lys Asn Asn Asn Ile Asn Gly Lys Gln Ile Asp Ala Lys Lys His
 435 440 445
 Ile Pro Trp Leu Ser Ile Gln Val Ile Val Phe Thr Thr Ser Ile Leu
 450 455 460
 Leu Gly Cys Ile Lys
 465

<210> 8

<211> 675

<212> DNA

<213> Ehrlichia canis

<220>

<221> CDS

<222> (1)..(675)

<223> Protein translated from nucleotides 4,121 through
4,795 (ORF of unknown function).

<400> 8

atg gta tta ttt atg aaa gct cat agc aca agt ata cgg aac ttt cag	48
Met Val Leu Phe Met Lys Ala His Ser Thr Ser Ile Arg Asn Phe Gln	
1 5 10 15	
cct tta gaa aga gct gct ata atc att gca gtg tta ggt tta gct gca	96
Pro Leu Glu Arg Ala Ala Ile Ile Ile Ala Val Leu Gly Leu Ala Ala	
20 25 30	
ttc ttg ttt gct gct gct gcc tgc agt gat cgt ttc caa aga ttg caa	144
Phe Leu Phe Ala Ala Ala Ala Cys Ser Asp Arg Phe Gln Arg Leu Gln	
35 40 45	
tta aca aat cca ttt gta ata gca gga atg gtt ggc ctt gca gtt ctt	192
Leu Thr Asn Pro Phe Val Ile Ala Gly Met Val Gly Leu Ala Val Leu	
50 55 60	
tta gtt gct tcc tta aca gca gca tta agt ata tgc tta act aaa agt	240
Leu Val Ala Ser Leu Thr Ala Ala Leu Ser Ile Cys Leu Thr Lys Ser	
65 70 75 80	
aag caa gtc aca caa cat gct att aga cat cgc ttt gga tac gag tca	288
Lys Gln Val Thr Gln His Ala Ile Arg His Arg Phe Gly Tyr Glu Ser	
85 90 95	
agc act tct tct tct gta ctg ctt gca ata tca ata att tct tta tta	336
Ser Thr Ser Ser Ser Val Leu Leu Ala Ile Ser Ile Ile Ser Leu Leu	
100 105 110	
ctt gct gca gca ttt tgt gga aag ata atg ggt aat gac aac cca gat	384
Leu Ala Ala Ala Phe Cys Gly Lys Ile Met Gly Asn Asp Asn Pro Asp	
115 120 125	
cta ttc ttt agc aag atg caa gaa ctc tcc aat cca ctt gtt gtt gca	432
Leu Phe Phe Ser Lys Met Gln Glu Leu Ser Asn Pro Leu Val Val Ala	
130 135 140	
gct att gta gcc gtt tct gtt ttc cta ctc tca ttc gta atg tat gct	480
Ala Ile Val Ala Val Ser Val Phe Leu Leu Ser Phe Val Met Tyr Ala	
145 150 155 160	
gca aag aac att ata agt cca gat aaa caa act cac gtt att ata tta	528
Ala Lys Asn Ile Ile Ser Pro Asp Lys Gln Thr His Val Ile Ile Leu	
165 170 175	

16

tct	aat	caa	caa	act	ata	gaa	gaa	gca	aaa	gta	gat	caa	gga	atg	aat	576
Ser	Asn	Gln	Gln	Thr	Ile	Glu	Glu	Ala	Lys	Val	Asp	Gln	Gly	Met	Asn	
			180					185					190			

att	ttg	tca	gca	gta	ctc	cca	gca	gct	ggc	att	gac	atc	atg	act	ata	624
Ile	Leu	Ser	Ala	Val	Leu	Pro	Ala	Ala	Gly	Ile	Asp	Ile	Met	Thr	Ile	
		195					200				205					

gct	tct	tgt	gac	att	tta	gca	gtg	agc	agc	cgg	gga	tcc	tct	cag	cat	672
Ala	Ser	Cys	Asp	Ile	Leu	Ala	Val	Ser	Ser	Arg	Gly	Ser	Ser	Gln	His	
	210					215					220					

caa																675
Gln																
225																

<210> 9

<211> 225

<212> PRT

<213> Ehrlichia canis

<400> 9

Met	Val	Leu	Phe	Met	Lys	Ala	His	Ser	Thr	Ser	Ile	Arg	Asn	Phe	Gln	
1				5					10					15		
Pro	Leu	Glu	Arg	Ala	Ala	Ile	Ile	Ile	Ala	Val	Leu	Gly	Leu	Ala	Ala	
			20					25					30			
Phe	Leu	Phe	Ala	Ala	Ala	Ala	Cys	Ser	Asp	Arg	Phe	Gln	Arg	Leu	Gln	
		35					40					45				
Leu	Thr	Asn	Pro	Phe	Val	Ile	Ala	Gly	Met	Val	Gly	Leu	Ala	Val	Leu	
	50					55					60					
Leu	Val	Ala	Ser	Leu	Thr	Ala	Ala	Leu	Ser	Ile	Cys	Leu	Thr	Lys	Ser	
	65				70					75				80		
Lys	Gln	Val	Thr	Gln	His	Ala	Ile	Arg	His	Arg	Phe	Gly	Tyr	Glu	Ser	
				85				90						95		
Ser	Thr	Ser	Ser	Ser	Val	Leu	Leu	Ala	Ile	Ser	Ile	Ile	Ser	Leu	Leu	
			100					105					110			
Leu	Ala	Ala	Ala	Phe	Cys	Gly	Lys	Ile	Met	Gly	Asn	Asp	Asn	Pro	Asp	
	115						120					125				
Leu	Phe	Phe	Ser	Lys	Met	Gln	Glu	Leu	Ser	Asn	Pro	Leu	Val	Val	Ala	
	130					135					140					
Ala	Ile	Val	Ala	Val	Ser	Val	Phe	Leu	Leu	Ser	Phe	Val	Met	Tyr	Ala	
145					150					155					160	

Ala	Lys	Asn	Ile	Ile	Ser	Pro	Asp	Lys	Gln	Thr	His	Val	Ile	Ile	Leu	
				165					170					175		
Ser	Asn	Gln	Gln	Thr	Ile	Glu	Glu	Ala	Lys	Val	Asp	Gln	Gly	Met	Asn	
				180					185					190		
Ile	Leu	Ser	Ala	Val	Leu	Pro	Ala	Ala	Gly	Ile	Asp	Ile	Met	Thr	Ile	
				195					200					205		
Ala	Ser	Cys	Asp	Ile	Leu	Ala	Val	Ser	Ser	Arg	Gly	Ser	Ser	Gln	His	
				210					215					220		
Gln																
225																

gat	cag	gta	agt	aaa	tgg	tat	gta	gta	aat	ttg	ata	gga	gat	aaa	ggt	48
Asp	Gln	Val	Ser	Lys	Trp	Tyr	Val	Val	Asn	Leu	Ile	Gly	Asp	Lys	Gly	
1				5					10					15		
gta	ata	gag	ata	tta	agc	ttc	ttg	cgc	ttt	act	aca	gtg	tgg	aat	gct	96
Val	Ile	Glu	Ile	Leu	Ser	Phe	Leu	Arg	Phe	Thr	Thr	Val	Trp	Asn	Ala	
			20					25					30			
gga	att	agt	ttt	ggt	ata	tta	aat	aac	ttt	gaa	tat	agt	aat	gtt	gtt	144
Gly	Ile	Ser	Phe	Gly	Ile	Leu	Asn	Asn	Phe	Glu	Tyr	Ser	Asn	Val	Val	
		35					40					45				
ttt	tgt	agt	atc	tcg	att	ttg	att	act	tgt	gtt	tta	tgc	tac	tta	ttt	192
Phe	Cys	Ser	Ile	Ser	Ile	Leu	Ile	Thr	Cys	Val	Leu	Cys	Tyr	Leu	Phe	
	50					55					60					
ata	gta	cag	cca	cat	tat	aga	tta	cct	ctt	gta	atc	att	att	ggg	ggg	240
Ile	Val	Gln	Pro	His	Tyr	Arg	Leu	Pro	Leu	Val	Ile	Ile	Ile	Gly	Gly	
65					70					75					80	

18

tca ata gga aat atc ata gat aga ata aga tat ggt gct gtc tat gat 288
 Ser Ile Gly Asn Ile Ile Asp Arg Ile Arg Tyr Gly Ala Val Tyr Asp
 85 90 95

ttt ata gat ttt tat atc aat aac tta cat tgg cct gta ttc aac ctg 336
 Phe Ile Asp Phe Tyr Ile Asn Asn Leu His Trp Pro Val Phe Asn Leu
 100 105 110

gcg gat tct ttt ata ttt tta ggt ata gta ata ata atg gca aag agt 384
 Ala Asp Ser Phe Ile Phe Leu Gly Ile Val Ile Ile Met Ala Lys Ser
 115 120 125

aat aac cac atg aaa caa att aac tgt aac tcc 417
 Asn Asn His Met Lys Gln Ile Asn Cys Asn Ser
 130 135

<210> 11

<211> 139

<212> PRT

<213> Ehrlichia canis

<400> 11

Asp Gln Val Ser Lys Trp Tyr Val Val Asn Leu Ile Gly Asp Lys Gly
 1 5 10 15

Val Ile Glu Ile Leu Ser Phe Leu Arg Phe Thr Thr Val Trp Asn Ala
 20 25 30

Gly Ile Ser Phe Gly Ile Leu Asn Asn Phe Glu Tyr Ser Asn Val Val
 35 40 45

Phe Cys Ser Ile Ser Ile Leu Ile Thr Cys Val Leu Cys Tyr Leu Phe
 50 55 60

Ile Val Gln Pro His Tyr Arg Leu Pro Leu Val Ile Ile Ile Gly Gly
 65 70 75 80

Ser Ile Gly Asn Ile Ile Asp Arg Ile Arg Tyr Gly Ala Val Tyr Asp
 85 90 95

Phe Ile Asp Phe Tyr Ile Asn Asn Leu His Trp Pro Val Phe Asn Leu
 100 105 110

Ala Asp Ser Phe Ile Phe Leu Gly Ile Val Ile Ile Met Ala Lys Ser
 115 120 125

Asn Asn His Met Lys Gln Ile Asn Cys Asn Ser
 130 135